



Antidepressants Help Men Decrease Alcohol Consumption, but Not Women

Libraries

Medical News

Keywords

ANTIDEPRESSANTS, CIHR, IRSC, ALCOHOL, DEPRESSION, WOMEN, MEN, CAMH

Contact Information

Available for logged-in reporters only

Description

CIHR-funded study explored the relationship between use of antidepressants and level of alcohol consumption, examining whether using antidepressants affected the link between depression and level of alcohol consumption. The research concluded that women suffering from depression consumed more alcohol than women who did not experience depression, regardless of antidepressant use. This finding differs significantly from rates found in male counterparts.

Newswise — CIHR-funded study explored the relationship between use of antidepressants and level of alcohol consumption, examining whether using antidepressants affected the link between depression and level of alcohol consumption. The research conducted by the Centre for Addiction and Mental Health (CAMH) concluded that women suffering from depression consumed more alcohol than women who did not experience depression, regardless of antidepressant use. This finding differs significantly from rates found in male counterparts. While men suffering from depression generally consume more alcohol than non-depressed men, those who use antidepressants consume alcohol at about the same level as non-depressed men.

Dr. Kathryn Graham, Senior Scientist with CAMH and Agnes Massak, Ph.D student in the Department of Psychology at the University of Western Ontario, published the study in the *Canadian Medical Association Journal (CMAJ)* on February 27, 2007.

“Our results agree with previous clinical research that suggests that the use of antidepressants is associated with lower alcohol consumption among men suffering from depression,” said Dr. Graham. “But this does not appear to be true for women.”

Overall, participants in the survey experiencing depression (both men and women) drank more alcohol than did non-depressed respondents. However, men taking antidepressants

consumed significantly less alcohol than depressed men who did not use antidepressants. Non-depressed men consumed 436 drinks per year, compared to 579 drinks for depressed men not using antidepressants, and 414 drinks for depressed men who used antidepressants.

Unfortunately for women, the alcohol use remained higher whether those experiencing depression took antidepressants or not. The numbers are telling: 179 drinks per year for non-depressed women, 235 drinks for depressed women not using antidepressants, and 264 drinks for depressed women who used antidepressants

“The fact that the relationship between the use of antidepressants and the level of alcohol consumption is different for men and women points to the importance of taking gender influences and sex differences into consideration in the treatment and prevention of many health conditions,” said Dr. Miriam Stewart, Scientific Director for CIHR Institute of Gender and Health. “This type of research reporting significant sex differences helps identify important clues for tailoring interventions.”

The scientists behind this study say further research is needed to assess whether this finding is due to drug effects or some other factor.

“We do not know whether antidepressants have different pharmacological effects on men and women, whether depression differs by gender, or whether the differences in the process of being treated for depression account for this discrepancy,” mentioned Dr. Graham. “For example, physicians prescribing antidepressants may be more likely to caution men than women about their drinking.”

For the study, 14,063 Canadian residents aged 18-76 years were surveyed. The survey included measures of quantity, frequency of drinking, depression and antidepressants use, over the period of a year.

The researchers used data from the GENACIS Canada survey, part of an international collaboration to investigate the influence of cultural variation on gender differences in alcohol use and related problems. CIHR provided over 1.3 million to GENACIS Canada (GENder Alcohol and Culture: an International Study). Over 35 countries and more than 100 leading alcohol and gender researchers are involved in the multinational study.

Every year, 12% of Canadians between the ages of 15 and 64 years suffer from some form of a mental disorder or substance dependence.

-30-

The Canadian Institutes of Health Research (CIHR) is the Government of Canada's agency for health research. CIHR's mission is to create new scientific knowledge and to catalyze its translation into improved health, more effective health services and products, and a strengthened Canadian health care system. Composed of 13 Institutes, CIHR provides leadership and support to more than 10,000 health researchers and trainees across Canada. www.cihr-irsc.gc.ca

The Centre for Addiction and Mental Health (CAMH) is one of the largest addiction and mental health organizations in North America and Canada's leading mental health and addiction teaching hospital. CAMH is a Pan American Health Organization and World Health Organization Collaborating Centre, and is fully affiliated with the University of Toronto. CAMH combines clinical care, research, policy, education and health promotion to improve the lives of people impacted by mental health and addiction issues.

www.camh.net

Ce document est également disponible en français.

[© 2007 Newswise](#). All Rights Reserved.