

Asian American and Pacific Islander (AA/PI) Community and Mental Health FACT SHEET

Did you know...

- There is very little research about mental health issues in these populations.
- Overall prevalence rates of diagnosable mental illnesses among AA/PIs are similar to those of the Caucasian population.
- Asian Americans and Pacific Islanders have the lowest rates of utilization of mental health services among ethnic populations.
- AA/PIs show higher levels of depressive symptoms than whites. South Asian refugees reported the highest rates among Asian groups.
- The Chinese American Psychiatric Epidemiologic Study (CAPES study) found a lifetime prevalence rate of about 7 percent (lifetime) and a 3 percent rate over one year.
- The National Comorbidity Study (NCS) reported even higher rates of major depression among Chinese Americans (17 percent for lifetime and 10 percent current).
- The word depression does not exit in certain Asian languages (e.g. Chinese).
- The suicide rates for Filipino (3.5 percent), Chinese (8.3 percent) and Japanese (9.1 percent) Americans are substantially lower than the suicide rate of white Americans (12.8 percent).
- Among elderly women of all ethnic or racial groups, Asians have the highest suicide rate.
- According to mental health care providers: Asian American women ages 15-24 have a higher rate of suicide than Caucasians, African Americans and Latinos in that age group.
- The Commonwealth Fund Survey of the Health of Adolescent Girls reported that Asian American adolescent girls had the highest rates of depressive symptoms of all racial/ethnic and gender groups.
- Southeast Asians suffer from particularly high rates of depression and post-traumatic stress disorder and exhibit more than twice the need for outpatient mental health services than the general Asian population.
- Suicide rates are higher than the national average for some groups of Asian Americans.
- The suicide rate among Asian Americans and Pacific Islanders in California is similar to that of the total population.
- Many Southeast Asian refugees are at risk for post-traumatic stress disorder (PTSD) associated with trauma experienced before and after immigration to the United States.
- One study found that 70 percent of Southeast Asian refugees receiving mental health care met diagnostic criteria for PTSD.
- In a study of Cambodian adolescents who survived Pol Pot's concentration camps, nearly half experienced PTSD and 41 percent suffered from depression 10 years after leaving Cambodia.
- Approximately 70 AA/PI providers are available for every 100,000 AA/PIs in the United States, compared to 173 per 100,000 Caucasians.
- AA/PIs appear to have the extremely low utilization of mental health services relative to other U.S. populations.
- In the CAPES study, only 17 percent of those experiencing problems sought care.
- A national study concluded that Asian Americans were one quarter as likely as whites to seek mental health services and half as likely as Latinos and African Americans. **

Created by the NAMI Multicultural & International Outreach Center June, 2003

References

- i Karen L. Koh, MPH and Margaret W. Leung, MPH. Asian Pacific Psychological Services, 431 30thStreet, Suite 6A, Oakland, CA 94609
- ii Mental Health: Culture, Race, Ethnicity Supplement to Mental Health: Report of the Surgeon General.
- iii Breaking the Silence. A Study of Depression Among Asian American Women. National Asian Women's Health Organization, 2001.
- iv Mental Health: Culture, Race, Ethnicity Supplement to Mental Health: Report of the Surgeon General.
- v The Surgeon Gereral's Call to Action to Prevent Suicide, 1999.
- vi Monthly Vital Statistics Report. Center for Disease Control and Prevention/National Center for Health Statistics; Vol. 46, No. 1. August 17, 1997 vii Louis Harris and Associates, Inc., 1997.
- viii Karen L. Koh, MPH and Margaret W. Leung, MPH. Asian Pacific Psychological Services, 431 30th Street, Suite 6A, Oakland, CA 94609
- ix The Surgeon Gereral's Call to Action to Prevent Suicide, 1999.
- x Mental Health: Culture, Race, Ethnicity Supplement to Mental Health: Report of the Surgeon General.
- xi Mental Health: Culture, Race, Ethnicity Supplement to Mental Health: Report of the Surgeon General.
- xii Mental Health: Culture, Race, Ethnicity Supplement to Mental Health: Report of the Surgeon General.
- xiii Snowden, LR. African American service use for mental health problems. Journal of Community Psychology.